

On Gossamer Metals and Insulating Behavior

George Chapline[□], Zaira Nazario[†] and David I. Santiago^{†,*}

[□] *Physics and Advanced Technologies Directorate*

Lawrence Livermore National Laboratory, Livermore, CA 94550.

[†] *Department of Physics, Stanford University, Stanford, California 94305*

^{*} *Gravity Probe B Relativity Mission, Stanford, California 94305*

We extend the Gossamer technique recently proposed to describe superconducting ground states to metallic ground states. The gossamer metal in a single band model will describe a metallic phase that becomes arbitrarily hard to differentiate from an insulator as one turns the Coulomb correlations up. We were motivated by the phase diagram of V_2O_3 and f-electron systems which have phase diagrams in which a line of first order metal-insulator transition ends at a critical point above which the two phases are indistinguishable. This means that one can go continuously from the metal to the “insulator”, suggesting that they might be the same phase.

PACS numbers: 71.27.+a, 71.10.-w, 71.30.+h, 71.15.-m

Vanadium sesquioxide V_2O_3 , one of the so called Mott insulators, has been widely studied because of its very interesting phase diagram¹. Like most Mott insulators, V_2O_3 is an antiferromagnet at low enough temperatures. This may be a general phenomenon and all correlated insulators will order at a low enough temperature.

The phase diagram of V_2O_3 and other Mott insulators is such that the antiferromagnetism and insulation disappears with pressure at $T = 0$ ¹. As one increases the temperature of the antiferromagnetic phase, the spin order melts and one is left with a spin disordered phase that is insulator-like in the sense that the resistivity is rising with diminishing temperature. As one applies pressure to this insulating-like phase it undergoes a first order phase transition into a metal.

The interesting thing is that the transition line between the metal and the insulator-like phase terminates at a critical point (T_c, P_c) . Similar behavior is observed in f-electron compounds⁵. This means that the metal and the insulating-like phase cannot be fundamentally different as one can go continuously from one into the other, analogous to the continuous transformation of liquid into a gas above the critical temperature. We posit that the spin disordered insulator-like phase is a bad tenuous “Gossamer” metal rather than a true insulator.

Recently Laughlin² and collaborators³ have advocated a similar Gossamer phenomenology as apt to describe the superconducting cuprates. The main idea of the Gossamer superconductor is that there is a superconducting pairing amplitude all the way to zero doping into the “insulating” regime. The Coulomb interactions decrease the superfluid density. The superconducting d-wave spectrum will be there all the time, but the spectral weight just becomes smaller and smaller as one goes to zero doping. The missing spectral weight goes into forming Hubbard-like bands. In the model introduced by Laughlin², this spectral redistribution comes about as result of partial Gutzwiller projection, which we call the Gossamer technique.

The Gossamer technique used for the superconductor should work for a metal in an analogous fashion, as has

been recently proposed⁴. Near full projection and near half-filling the density of states at the Fermi level collapses to zero. The missing spectral weight would go to forming Mott-Hubbard bands. Such a “Gossamer” metal would be impossible to tell apart from an insulator except at the lowest possible temperatures in which, at least for the case of V_2O_3 , antiferromagnetic order intervenes and the material is truly insulating. If antiferromagnetism does not intervene at low temperatures, we would expect the resistivity to saturate to a large but finite value at small temperatures absent localization effects. We propose that something analogous to this is happening for the disordered insulating-like phase of some correlated electron systems.

We speculate that, while our original motivation for the present work was V_2O_3 , perhaps f-electron systems, which are charge transfer insulators like the superconducting cuprates, are more apt to be described as gossamer metals. In particular we draw attention to f-electron collapse in the rare earth and actinide elements and their compounds⁶. This is usually a first order phase transition with a volume change, and results from changes in composition or pressure. It stands out as a characteristic feature of these materials. Indeed understanding f-electron collapse may be of importance for understanding the remarkable properties of these materials. A typical example of f-electron collapse is the $\alpha \rightarrow \gamma$ phases transition in elemental cerium, which occurs at room temperature at a pressure near to 10 kbar⁵. This phase transition is thought to be the result of an f-electron initially localized on a cerium ion becoming delocalized as a result of increased hybridization with conduction electrons. Correlations between the f-electrons are also thought to play an important role. Although f-electron collapse is normally a first order phase transition, in the following we would like to focus on the fact that since there is a critical point the α and γ phases of Ce can really be regarded as a single Gossamer metallic phase.

The Gossamer technique is a bit more delicate to apply to a metallic Fermi sea ground state than to a supercon-

ducting BCS ground state. The reason is that the abundance of low energy degrees of freedom in the metal might make the projection uncontrolled in the infrared leading to unphysical results. Since the superconductor does not have such a plethora of low energy degrees of freedom (irrespective of whether the gap has nodes or not), the calculation has no infrared problems: it is regularized by the superconducting order. We therefore will borrow heavily from the calculations for the superconductor in previous work^{2,3}. After obtaining the results, we collapse the gap to zero to study the physics of Gossamer metals.

We now proceed to reproduce and review the previous results^{2,3} and apply them to describe bad metallic behavior. The Gossamer ground states are constructed by applying the partial Gutzwiller “projector”

$$\Pi_{\alpha_0} = \prod_j z_0^{(n_{j\uparrow} + n_{j\downarrow})/2} (1 - \alpha_0 n_{j\uparrow} n_{j\downarrow}) \quad (1)$$

$0 \leq \alpha_0 < 1$ is a measure of how effective the projector is and in a real material it will be related to the Coulomb repulsion. The factor of z_0 , the quantum fugacity, in the projector is the extra probability of having an electron at site j *chosen* to keep the total number of particles constant at $(1 - \delta)N$ after projecting. The fugacity is given by $z_0 = (\sqrt{1 - \alpha(1 - \delta^2)} - \delta) / [(1 - \alpha)(1 - \delta)]$ with $(1 - \alpha_0)^2 = 1 - \alpha^2$.

The Gossamer superconducting ground state is postulated to be $|\Psi\rangle = \Pi_{\alpha} |\Phi\rangle$. Here $|\Phi\rangle$ is the BCS ground state:

$$|\Phi\rangle = \prod_{\vec{k}} (u_{\vec{k}} + v_{\vec{k}} c_{\vec{k}\uparrow}^{\dagger} c_{-\vec{k}\downarrow}^{\dagger}) |0\rangle \quad (2)$$

where $u_{\vec{k}}, v_{\vec{k}}$ are the BCS coherence factors given by $u_{\vec{k}} = \sqrt{(E_{\vec{k}} + \epsilon_{\vec{k}} - \mu)/2E_{\vec{k}}}$ and $v_{\vec{k}} = \sqrt{[E_{\vec{k}} - (\epsilon_{\vec{k}} - \mu)]/2E_{\vec{k}}}$ with dispersion $E_{\vec{k}} = \sqrt{(\epsilon_{\vec{k}} - \mu)^2 + \Delta_{\vec{k}}^2}$ where $\epsilon_{\vec{k}}$ is the kinetic energy of the metal measured from the Fermi level, μ is the chemical potential and $\Delta_{\vec{k}}$ is the superconducting gap. Such projected ground states have been studied before^{7,8,9,10}.

We never fully project ($\alpha_0 < 1$) in order for the partial projector to have an inverse:

$$\Pi_{\alpha}^{-1} = \prod_j z_0^{-(n_{j\uparrow} + n_{j\downarrow})/2} (1 + \beta_0 n_{j\uparrow} n_{j\downarrow}) \quad (3)$$

with $\beta_0 = \alpha_0/(1 - \alpha_0)$. The Gossamer ground state is the *exact* ground state of the Gossamer Hamiltonian:

$$\mathcal{H} = \sum_{\vec{k}\sigma} E_{\vec{k}} B_{\vec{k}\sigma}^{\dagger} B_{\vec{k}\sigma}, \quad B_{\vec{k}\sigma} |\Psi\rangle = 0. \quad (4)$$

where:

$$B_{\vec{k}\uparrow\{\downarrow\}} = \Pi_{\alpha} b_{\vec{k}\uparrow\{\downarrow\}} \Pi_{\alpha}^{-1} = \frac{1}{\sqrt{N}} \sum_j e^{i\vec{k}\cdot\vec{r}_j} \times \left[z_0^{-1/2} u_{\vec{k}} (1 + \beta_0 n_{j\downarrow\{\uparrow\}}) c_{j\uparrow\{\downarrow\}} \pm z_0^{1/2} v_{\vec{k}} (1 - \alpha_0 n_{j\uparrow\{\downarrow\}}) c_{j\downarrow\{\uparrow\}}^{\dagger} \right] \quad (5)$$

with $b_{\vec{k}\uparrow\{\downarrow\}} = u_{\vec{k}} c_{j\uparrow\{\downarrow\}} \pm v_{\vec{k}} c_{j\downarrow\{\uparrow\}}^{\dagger}$ the Bogoliubov quasiparticle operators which annihilate the BCS ground state. The Gossamer Hamiltonian would be impossible to define if we used a full Gutzwiller projector.

The Gossamer ground state is adiabatically continuable to the BCS ground state by continuously varying α_0 to zero. Its uniqueness follows from the uniqueness of the BCS ground state up to a phase. Therefore the Gossamer superconductor describes the same phase of matter as the BCS superconductor. In a similar fashion once we collapse the gap to obtain the Gossamer metal ground state, it will be adiabatically deformable to a regular Fermi sea metallic ground state and, hence they will be the same zero temperature phase of matter.

In previous work² it was shown that the variational wavefunction $|\vec{k}\sigma\rangle = \Pi_{\alpha} b_{\vec{k}\sigma}^{\dagger} |\Phi\rangle$ represents an appropriate approximation to the low energy quasiparticle excitations and that, remarkably, their dispersion does not change under projection:

$$\frac{\langle \vec{k}\sigma | \mathcal{H} | \vec{k}\sigma \rangle}{\langle \vec{k}\sigma | \vec{k}\sigma \rangle} = E_{\vec{k}} \frac{\langle \Psi | \Psi \rangle}{\langle \vec{k}\sigma | \vec{k}\sigma \rangle} \simeq E_{\vec{k}} \quad (6)$$

We collapse the superconducting gap to find that the dispersion in the the Gossamer metal is *unchanged* from the dispersion in the regular metal. This might be a bit surprising as usually correlation effects are thought to make the charge carriers heavy¹¹. What is happening here is not that the carriers are getting arbitrarily heavy as we approach the transition, but the metallic band is becoming thinner and the missing spectral weight goes to energies far from the Fermi sea to forming Hubbard bands. The carriers are just as fast, there are just less of them which degrades the conductivity.

From the matrix elements²

$$\begin{aligned} & \frac{\langle \Phi | c_{j\uparrow} \Pi_{\alpha}^2 c_{j'\uparrow}^{\dagger} | \Phi \rangle}{\langle \Phi | \Pi_{\alpha}^2 | \Phi \rangle} \times \frac{\langle \Phi | \Phi \rangle}{\langle \Phi | c_{j\uparrow} c_{j'\uparrow}^{\dagger} | \Phi \rangle} \\ & \simeq \frac{4}{1 - \delta^2} (z z_0) \left[\frac{1 + (1 - \alpha)z}{1 + 2z + (1 - \alpha)z^2} \right]^2 = 1 \end{aligned} \quad (7)$$

and

$$\frac{\langle \Phi | c_{j\uparrow}^{\dagger} \Pi_{\alpha}^2 c_{j'\uparrow} | \Phi \rangle}{\langle \Phi | \Pi_{\alpha}^2 | \Phi \rangle} \times \frac{\langle \Phi | \Phi \rangle}{\langle \Phi | c_{j\uparrow}^{\dagger} c_{j'\uparrow} | \Phi \rangle}$$

$$\simeq \frac{4}{1-\delta^2} \left(\frac{z}{z_0} \right) \left[\frac{1+z}{1+2z+(1-\alpha)z^2} \right]^2 = 1 \quad (8)$$

the photoemission amplitudes were calculated for the Gossamer superconductor. After collapsing the gap, for the Gossamer metal these are given by

$$\frac{\langle -\vec{k} \downarrow | c_{\vec{k}\uparrow}^\dagger | \Psi \rangle}{\sqrt{\langle -\vec{k} \downarrow | -\vec{k} \downarrow \rangle \langle \Psi | \Psi \rangle}} = 0 \quad ,$$

$$\frac{\langle -\vec{k} \downarrow | c_{\vec{k}\uparrow}^\dagger | \Psi \rangle}{\sqrt{\langle -\vec{k} \downarrow | -\vec{k} \downarrow \rangle \langle \Psi | \Psi \rangle}} = g \quad , \quad (9)$$

with

$$g^2 \simeq \frac{2\alpha_0}{\alpha} \left\{ 1 - \frac{\alpha_0}{\alpha} \left[\frac{1 - \sqrt{1 - \alpha(1 - \delta^2)}}{1 - \delta^2} \right] \right\} \quad . \quad (10)$$

The suppression of photoemission amplitude is endemic of the smaller number of metallic electrons whose number is diminished from that in the unprojected metal by a factor g^2 which goes to $2|\delta|/(1+|\delta|)$ as $\alpha_0 \rightarrow 1$. This is consistent with the previous result² that the superfluid density in the Gossamer superconductor goes like g^2 as

there is a sum rule for the superconductor making the density of conducting electrons equal to the superfluid density. For strong correlations the number of metallic electrons vanishes at half-filling. The material is metallic at all other dopings. Whether there is a full gap opened at zero doping is an interesting question. We have not yet been able to determine if the whole Fermi surface is destroyed or if there are some Fermi points as the material is on the brink of becoming insulating and opening a gap. The continuous behavior with doping certainly argues for the latter.

We now proceed to analyze the specific form that the Gossamer metal Hamiltonian takes. After some manipulation, we can bring the Hamiltonian into the form

$$\mathcal{H} = \sum_{\vec{k}} \frac{E_{\vec{k}}}{N} \sum_{i,j}^N e^{-i\vec{k}(\vec{r}_i - \vec{r}_j)} \{ z_0^{-1} (1 + \beta_0 n_{i\downarrow}) (1 + \beta_0 n_{j\downarrow}) c_{i\uparrow}^\dagger c_{j\uparrow} + z_0^{-1} (1 + \beta_0 n_{i\uparrow}) (1 + \beta_0 n_{j\uparrow}) c_{i\downarrow}^\dagger c_{j\downarrow} \} \quad (11)$$

We will see that the sums break up into a chemical potential, kinetic, as well as a Hubbard U terms.

Kinetic Term in the Gossamer Metal: The off-site contributions of $\mathcal{H}$ give a hopping (kinetic) term in the Hamiltonian. We can write $\mathcal{H}$ as a sum of on-site and off-site contributions, $\mathcal{H} = \mathcal{H}_{on\ site} + \mathcal{H}_{off\ site}$, where the two contributions read:

$$\mathcal{H}_{on\ site} = \sum_{\vec{k}} \frac{E_{\vec{k}}}{N} \sum_j^N \{ z_0^{-1} (1 + \beta_0 n_{j\downarrow}) (1 + \beta_0 n_{j\downarrow}) c_{j\uparrow}^\dagger c_{j\uparrow} + \{ \uparrow \rightleftharpoons \downarrow \} \} \quad (12)$$

$$\mathcal{H}_{off\ site} = \sum_{\vec{k}} \frac{E_{\vec{k}}}{N} \sum_{i \neq j}^N e^{-i\vec{k}(\vec{r}_i - \vec{r}_j)} \times \{ z_0^{-1} (1 + \beta_0 n_{i\downarrow}) (1 + \beta_0 n_{j\downarrow}) c_{i\uparrow}^\dagger c_{j\uparrow} + \{ \uparrow \rightleftharpoons \downarrow \} \} \quad (13)$$

In this section we are interested only on the off-site contribution. Without the partial Gutzwiller projection ($\alpha_0 = 0$, $z_0 = 1$) this term is just the kinetic term of a mean-field Hamiltonian $\sum_{\vec{k}\sigma} (\epsilon_{\vec{k}} - \mu) c_{\vec{k}\sigma}^\dagger c_{\vec{k}\sigma}$. However, the partial projection induces complications which make $\mathcal{H}_{off\ site}$ very difficult to diagonalize analitically. In order to get a rough estimate for the change in the kinetic term due to partial projection, we will make the mean field approximation $\langle n_{i\uparrow} \rangle = \langle n_{i\downarrow} \rangle = \frac{1}{2}(1 - \delta)$, and we will replace the number operators with this average. The

term then becomes:

$$\mathcal{H}_{off\ site} = \sum_{\vec{k}\sigma} \frac{E_{\vec{k}}}{N} \sum_{i \neq j}^N e^{-i\vec{k}(\vec{r}_i - \vec{r}_j)} z_0^{-1} \left(1 + \frac{\beta_0(1 - \delta)}{2} \right)^2 c_{i\sigma}^\dagger c_{j\sigma} \quad (14)$$

This term finally is:

$$\mathcal{H}_{off\ site} = z_0^{-1} \left(1 + \frac{\beta_0(1 - \delta)}{2} \right)^2 \sum_{\vec{k}\sigma} (\epsilon_{\vec{k}} - \mu) c_{\vec{k}\sigma}^\dagger c_{\vec{k}\sigma} \quad (15)$$

Therefore the effect of the partial projection on the kinetic term in the Gossamer Hamiltonian is an overall

constant, a renormalization of the energy by

$$\frac{[2 - \alpha_0(1 + \delta)]^2(1 - \delta)}{4(\sqrt{1 - \alpha(1 - \delta^2)} - \delta)}, \quad (16)$$

with $1 - \alpha = (1 - \alpha_0)^2$. As we can see, the constant blows up as one approaches full projection $\alpha_0 \rightarrow 1$. This does not lead to any complications since as we will soon see, the other terms blow up in an exactly similar fashion leaving their physically relevant ratio finite. When unprojected the renormalization factor is 1 as it should be.

Hubbard-U Term in Gossamer Metal: The Hubbard-U term will arise out of the on-site contribution, $\mathcal{H}_{on\ site}$ which was explicitly written down in the previous section. One can write the terms as:

$$\begin{aligned} \mathcal{H}_{on\ site} = \sum_{\vec{k}} \frac{E_{\vec{k}}}{N} \sum_j^N \{ z_0^{-1}(n_{j\uparrow} + n_{j\downarrow}) + \\ + z_0^{-1}(4\beta_0 + 2\beta_0^2)n_{j\uparrow}n_{j\downarrow} \} \end{aligned} \quad (17)$$

The first term is a chemical potential, and the second term is the Hubbard-U. We will concern ourselves only with this last term. For the factor containing β_0 we obtain:

$$z_0^{-1}(4\beta_0 + 2\beta_0^2) = \frac{2\alpha_0(2 - \alpha_0)(1 - \delta)}{\sqrt{1 - \alpha(1 - \delta^2)} - \delta} \quad (18)$$

and therefore the Gossamer metal will have a Hubbard U term of the form $\sum_j^N U n_{j\uparrow} n_{j\downarrow}$ where, at half-filling

$$U = \sum_{\vec{k}} \frac{E_{\vec{k}}}{N} \frac{2\alpha_0(2 - \alpha_0)(1 - \delta)}{\sqrt{1 - \alpha(1 - \delta^2)} - \delta} \quad (19)$$

As we can see, at full projection $\alpha_0 \rightarrow 1$ this term blows up as well, but the ratio between the hopping amplitude computed in the previous section and the Hubbard U remains finite and is of order 1 near full projection. When unprojected, the U term is zero as it should be.

The existence of the growing Hubbard U term means that as we go to half-filling and full projection magnetic correlations will get enhanced leading to a diverging magnetic susceptibility in the exact same way as for the Gossamer superconductor³ after we collapse the gap. The spectral weight will consist of Mott-Hubbard bands at high energies with the chemical potential pinned at midgap at an ever fainter band from which the Gossamer quasiparticles are excited with a dispersion unchanged from the noninteracting metal.

As mentioned before, this unmodified dispersion is different behavior from what was found in previous investigations of the Hubbard model with Gutzwiller projectors by Brinkman and Rice¹¹. The difference stems from the fact that the projection technique used in the previous work since it did not include a fugacity factor, z_0 , to conserve particle number. This leads to an arbitrarily large U and finite t leading to a diverging carrier mass as one goes to half-filling. In fact, a recent proposal for a Gossamer metal ground state⁴ with projectors that do not conserve the number of particles finds the exact same behavior as in the Brinkman-Rice work¹¹. In this note we make sure to conserve particle number, the ratio U/t remains finite and below the critical value for true insulation, and the carriers cannot become arbitrarily heavy.

In the present work we extended the Gossamer technique originally developed for the superconducting cuprates to metallic ground states. The Gossamer metal describes strongly correlated bad metal behavior that is very hard to distinguish from true insulating behavior, the implication being that the magnetically disordered insulating-like phases in some systems might really be a Gossamer metal with very much degraded conductivity. The degraded conductivity arises from a depletion of spectral weight of metallic electrons and not by an ever growing effective mass of the carriers. We studied this in a single band model. The single band model might not be an apt description of the f-electron system, which was one of the motivating players for the present work. Despite this, we believe that the general features we have described are general and should survive in such systems.

Acknowledgements: We thank Bob Laughlin for valuable discussions and suggestions. We thank B. A. Bernevig and D. Giuliano for discussions.

¹ D. B. McWhan, *et. al.* Phys. Rev. B **7**, 1920 (1973).

² R. B. Laughlin e-print arxiv:cond-mat/0209269 (2002).

³ B. A. Bernevig, R. B. Laughlin, and D. I. Santiago, Physical Review Letters in press; e-print arxiv:cond-mat/0303045(2003).

⁴ M. Kollar, e-print arxiv:cond-mat/0308513 (2003).

⁵ D. Koskimaki and K. Gschneider, Jr., Handbook on the Physics and Chemistry of the Rare Earths, , p. 337 edited

by K. A. Gschneider, Jr. and L. Eyring (North-Holland, Amsterdam, 1979).

⁶ U. Benedict, J. Alloys Compd. **193**, 88 (1993).

⁷ A. Paramekanti, *et. al.* Physical Review Letters **87** 21002 (2001).

⁸ F. Zhang, *et. al.*, Supercond. Sci & Tech **1**, 36 (1988).

⁹ F. C. Zhang e-print arxiv:cond-mat/0209272 (2002).

¹⁰ F. C. Zhang e-print arxiv:cond-mat/0308398 (2002).

- ¹¹ W. F. Brinkman and T. M. Rice, Phys. Rev. B **2**, 4302 (1970).